

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100 101 102 103 104 105 106 107 108 109 110 111 112 113 114 115 116 117 118 119 120 121 122 123 124 125 126 127 128 129 130 131 132 133 134 135 136 137 138 139 140 141 142 143 144 145 146 147 148 149 150 151 152 153 154 155 156 157 158 159 160 161 162 163 164 165 166 167 168 169 170 171 172 173 174 175 176 177 178 179 180 181 182 183 184 185 186 187 188 189 190 191 192 193 194 195 196 197 198 199 200 201 202 203 204 205 206 207 208 209 210 211 212 213 214 215 216 217 218 219 220 221 222 223 224 225 226 227 228 229 230 231 232 233 234 235 236 237 238 239 240 241 242 243 244 245 246 247 248 249 250 251 252 253 254 255 256 257 258 259 260 261 262 263 264 265 266 267 268 269 270 271 272 273 274 275 276 277 278 279 280 281 282 283 284 285 286 287 288 289 290 291 292 293 294 295 296 297 298 299 300 301 302 303 304 305 306 307 308 309 310 311 312 313 314 315 316 317 318 319 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 409 410 411 412 413 414 415 416 417 418 419 420 421 422 423 424 425 426 427 428 429 430 431 432 433 434 435 436 437 438 439 440 441 442 443 444 445 446 447 448 449 450 451 452 453 454 455 456 457 458 459 460 461 462 463 464 465 466 467 468 469 470 471 472 473 474 475 476 477 478 479 480 481 482 483 484 485 486 487 488 489 490 491 492 493 494 495 496 497 498 499 500 501 502 503 504 505 506 507 508 509 510 511 512 513 514 515 516 517 518 519 520 521 522 523 524 525 526 527 528 529 530 531 532 533 534 535 536 537 538 539 540 541 542 543 544 545 546 547 548 549 550 551 552 553 554 555 556 557 558 559 560 561 562 563 564 565 566 567 568 569 570 571 572 573 574 575 576 577 578 579 580 581 582 583 584 585 586 587 588 589 590 591 592 593 594 595 596 597 598 599 600 601 602 603 604 605 606 607 608 609 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 648 649 650 651 652 653 654 655 656 657 658 659 660 661 662 663 664 665 666 667 668 669 670 671 672 673 674 675 676 677 678 679 680 681 682 683 684 685 686 687 688 689 690 691 692 693 694 695 696 697 698 699 700 701 702 703 704 705 706 707 708 709 710 711 712 713 714 715 716 717 718 719 720 721 722 723 724 725 726 727 728 729 730 731 732 733 734 735 736 737 738 739 740 741 742 743 744 745 746 747 748 749 750 751 752 753 754 755 756 757 758 759 760 761 762 763 764 765 766 767 768 769 770 771 772 773 774 775 776 777 778 779 780 781 782 783 784 785 786 787 788 789 790 791 792 793 794 795 796 797 798 799 800 801 802 803 804 805 806 807 808 809 810 811 812 813 814 815 816 817 818 819 820 821 822 823 824 825 826 827 828 829 830 831 832 833 834 835 836 837 838 839 840 841 842 843 844 845 846 847 848 849 850 851 852 853 854 855 856 857 858 859 860 861 862 863 864 865 866 867 868 869 870 871 872 873 874 875 876 877 878 879 880 881 882 883 884 885 886 887 888 889 890 891 892 893 894 895 896 897 898 899 900 901 902 903 904 905 906 907 908 909 910 911 912 913 914 915 916 917 918 919 920 921 922 923 924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939 940 941 942 943 944 945 946 947 948 949 950 951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 967 968 969 970 971 972 973 974 975 976 977 978 979 980 981 982 983 984 985 986 987 988 989 990 991 992 993 994 995 996 997 998 999 1000 1001 1002 1003 1004 1005 1006 1007 1008 1009 1010 1011 1012 1013 1014 1015 1016 1017 1018 1019 1020 1021 1022 1023 1024 1025 1026 1027 1028 1029 1030 1031 1032 1033 1034 1035 1036 1037 1038 1039 1040 1041 1042 1043 1044 1045 1046 1047 1048 1049 1050 1051 1052 1053 1054 1055 1056 1057 1058 1059 1060 1061 1062 1063 1064 1065 1066 1067 1068 1069 1070 1071 1072 1073 1074 1075 1076 1077 1078 1079 1080 1081 1082 1083 1084 1085 1086 1087 1088 1089 1090 1091 1092 1093 1094 1095 1096 1097 1098 1099 1100 1101 1102 1103 1104 1105 1106 1107 1108 1109 1110 1111 1112 1113 1114 1115 1116 1117 1118 1119 1120 1121 1122 1123 1124 1125 1126 1127 1128 1129 1130 1131 1132 1133 1134 1135 1136 1137 1138 1139 1140 1141 1142 1143 1144 1145 1146 1147 1148 1149 1150 1151 1152 1153 1154 1155 1156 1157 1158 1159 1160 1161 1162 1163 1164 1165 1166 1167 1168 1169 1170 1171 1172 1173 1174 1175 1176 1177 1178 1179 1180 1181 1182 1183 1184 1185 1186 1187 1188 1189 1190 1191 1192 1193 1194 1195 1196 1197 1198 1199 1200 1201 1202 1203 1204 1205 1206 1207 1208 1209 1210 1211 1212 1213 1214 1215 1216 1217 1218 1219 1220 1221 1222 1223 1224 1225 1226 1227 1228 1229 1230 1231 1232 1233 1234 1235 1236 1237 1238 1239 1240 1241 1242 1243 1244 1245 1246 1247 1248 1249 1250 1251 1252 1253 1254 1255 1256 1257 1258 1259 1260 1261 1262 1263 1264 1265 1266 1267 1268 1269 1270 1271 1272 1273 1274 1275 1276 1277 1278 1279 1280 1281 1282 1283 1284 1285 1286 1287 1288 1289 1290 1291 1292 1293 1294 1295 1296 1297 1298 1299 1300 1301 1302 1303 1304 1305 1306 1307 1308 1309 1310 1311 1312 1313 1314 1315 1316 1317 1318 1319 1320 1321 1322 1323 1324 1325 1326 1327 1328 1329 1330 1331 1332 1333 1334 1335 1336 1337 1338 1339 1340 1341 1342 1343 1344 1345 1346 1347 1348 1349 1350 1351 1352 1353 1354 1355 1356 1357 1358 1359 1360 1361 1362 1363 1364 1365 1366 1367 1368 1369 1370 1371 1372 1373 1374 1375 1376 1377 1378 1379 1380 1381 1382 1383 1384 1385 1386 1387 1388 1389 1390 1391 1392 1393 1394 1395 1396 1397 1398 1399 1400 1401 1402 1403 1404 1405 1406 1407 1408 1409 1410 1411 1412 1413 1414 1415 1416 1417 1418 1419 1420 1421 1422 1423 1424 1425 1426 1427 1428 1429 1430 1431 1432 1433 1434 1435 1436 1437 1438 1439 1440 1441 1442 1443 1444 1445 1446 1447 1448 1449 1450 1451 1452 1453 1454 1455 1456 1457 1458 1459 1460 1461 1462 1463 1464 1465 1466 1467 1468 1469 1470 1471 1472 1473 1474 1475 1476 1477 1478 1479 1480 1481 1482 1483 1484 1485 1486 1487 1488 1489 1490 1491 1492 1493 1494 1495 1496 1497 1498 1499 1500 1501 1502 1503 1504 1505 1506 1507 1508 1509 1510 1511 1512 1513 1514 1515 1516 1517 1518 1519 1520 1521 1522 1523 1524 1525 1526 1527 1528 1529 1530 1531 1532 1533 1534 1535 1536 1537 1538 1539 1540 1541 1542 1543 1544 1545 1546 1547 1548 1549 1550 1551 1552 1553 1554 1555 1556 1557 1558 1559 1560 1561 1562 1563 1564 1565 1566 1567 1568 1569 1570 1571 1572 1573 1574 1575 1576 1577 1578 1579 1580 1581 1582 1583 1584 1585 1586 1587 1588 1589 1590 1591 1592 1593 1594 1595 1596 1597 1598 1599 1600 1601 1602 1603 1604 1605 1606 1607 1608 1609 1610 1611 1612 1613 1614 1615 1616 1617 1618 1619 1620 1621 1622 1623 1624 1625 1626 1627 1628 1629 1630 1631 1632 1633 1634 1635 1636 1637 1638 1639 1640 1641 1642 1643 1644 1645 1646 1647 1648 1649 1650 1651 1652 1653 1654 1655 1656 1657 1658 1659 1660 1661 1662 1663 1664 1665 1666 1667 1668 1669 1670 1671 1672 1673 1674 1675 1676 1677 1678 1679 1680 1681 1682 1683 1684 1685 1686 1687 1688 1689 1690 1691 1692 1693 1694 1695 1696 1697 1698 1699 1700 1701 1702 1703 1704 1705 1706 1707 1708 1709 1710 1711 1712 1713 1714 1715 1716 1717 1718 1719 1720 1721 1722 1723 1724 1725 1726 1727 1728 1729 1730 1731 1732 1733 1734 1735 1736 1737 1738 1739 1740 1741 1742 1743 1744 1745 1746 1747 1748 1749 1750 1751 1752 1753 1754 1755 1756 1757 1758 1759 1760 1761 1762 1763 1764 1765 1766 1767 1768 1769 1770 1771 1772 1773 1774 1775 1776 1777 1778 1779 1780 1781 1782 1783 1784 1785 1786 1787 1788 1789 1790 1791 1792 1793 1794 1795 1796 1797 1798 1799 1800 1801 1802 1803 1804 1805 1806 1807 1808 1809 1810 1811 1812 1813 1814 1815 1816 1817 1818 1819 1820 1821 1822 1823 1824 1825 1826 1827 1828 1829 1830 1831 1832 1833 1834 1835 1836 1837 1838 1839 1840 1841 1842 1843 1844 1845 1846 1847 1848 1849 1850 1851 1852 1853 1854 1855 1856 1857 1858 1859 1860 1861 1862 1863 1864 1865 1866 1867 1868 1869 1870 1871 1872 1873 1874 1875 1876 1877 1878 1879 1880 1881 1882 1883 1884 1885 1886 1887 1888 1889 1890 1891 1892 1893 1894 1895 1896 1897 1898 1899 1900 1901 1902 1903 1904 1905 1906 1907 1908 1909 1910 1911 1912 1913 1914 1915 1916 1917 1918 1919 1920 1921 1922 1923 1924 1925 1926 1927 1928 1929 1930 1931 1932 1933 1934 1935 1936 1937 1938 1939 1940 1941 1942 1943 1944 1945 1946 1947 1948 1949 1950 1951 1952 1953 1954 1955 1956 1957 1958 1959 1960 1961 1962 1963 1964 1965 1966 1967 1968 1969 1970 1971 1972 1973 1974 1975 1976 1977 1978 1979 1980 1981 1982 1983 1984 1985 1986 1987 1988 1989 1990 1991 1992 1993 1994 1995 1996 1997 1998 1999 2000 2001 2002 2003 2004 2005 2006 2007 2008 2009 2010 2011 2012 2013 2014 2015 2016 2017 2018 2019 2020 2021 2022 2023 2024 2025 2026 2027 2028 2029 2030 2031 2032 2033 2034 2035 2036 2037 2038 2039 2040 2041 2042 2043 2044 2045 2046 2047 2048 2049 2050 2051 2052 2053 2054 2055 2056 2057 2058 2059 2060 2061 2062 2063 2064 2065 2066 2067 2068 2069 2070 2071 2072 2073 2074 2075 2076 2077 2078 2079 2080 2081 2082 2083 2084 2085 2086 2087 2088 2089 2090 2091 2092 2093 2094 2095 2096 2097 2098 2099 2100 2101 2102 2103 2104 2105 2106 2107 2108 2109 2110 2111 2112 2113 2114 2115 2116 2117 2118 2119 2120 2121 2122 2123 2124 2125 2126 2127 2128 2129 2130 2131 2132 2133 2134 2135 2136 2137 2138 2139 2140 2141 2142 2143 2144 2145 2146 2147 2148 2149 2150 2151 2152 2153 2154 2155 2156 2157 2158 2159 2160 2161 2162 2163 2164 2165 2166 2167 2168 2169 2170 2171 2172 2173 2174 2175 2176 2177 2178 2179 2180 2181 2182 2183 2184 2185 2186 2187 2188 2189 2190 2191 2192 2193 2194 2195 2196 2197 2198 2199 2200 2201 2202 2203 2204 2205 2206 2207 2208 2209 2210 2211 2212 2213 2214 2215 2216 2217 2218 2219 2220 2221 2222 2223 2224 2225 2226 2227 2228 2229 2230 2231 2232 2233 2234 2235 2236 2237 2238 2239 2240 2241 2242 2243 2244 2245 2246 2247 2248 2249 2250 2251 2252 2253 2254 2255 2256 2257 2258 2259 2260 2261 2262 2263 2264 2265 2266 2267 2268 2269 2270 2271 2272 2273 2274 2275 2276 2277 2278 2279 2280 2281 2282 2283 2284 2285 2286 2287 2288 2289 2290 2291 2292 2293 2294 2295 2296 2297 2298 2299 2300 2301 2302 2303 2304 2305 2306 2307 2308 2309 2310 2311 2312 2313 2314 2315 2316 2317 2318 2319 2320 2321 2322 2323 2324 2325 2326 2327 2328 2329 2330 2331 2332 2333 2334 2335 2336 2337 2338 2339 2340 2341 2342 2343 2344 2345 2346 2347 2348 2349 2350 2351 2352 2353 2354 2355 2356 2357 2358 2359 2360 2361 2362 2363 2364 2365 2366 2367 2368 2369 2370 2371 2372 2373 2374 2375 2376 2377 2378 2379 2380 2381 2382 2383 2384 2385 2386 2387 2388 2389 2390 2391 2392 2393 2394 2395 2396 2397 2398 2399 2400 2401 2402 2403 2404 2405 2406 2407 2408 2409 2410 2411 2412 2413 2414 2415 2416 2417 2418 2419 2420 2421 2422 2423 2424 2425 2426 2427 2428 2429 2430 2431 2432 2433 2434 2435 2436 2437 2438 2439 2440 2441 2442 2443 2444 2445 2446 2447 2448 2449 2450 2451 2452 2453 2454 2455 2456 2457 2458 2459 2460 2461 2462 2463 2464 2465 2466 2467 2468 2469 2470 2471 2472 2473 2474 2475 2476 2477 2478 2479 2480 2481 2482 2483 2484 2485 2486 2487 2488 2489 2490 2491 2492 2493 2494 2495 2496 2497 2498 2499 2500 2501 2502 2503 2504 2505 2506 2507 2508 2509 2510 2511 2512 2513 2514 2515 2516 2517 2518 2519 2520 2521 2522 2523 2524 2525 2526 2527 2528 2529 2530 2531 2532 2533 2534 2535 2536 2537 2538 2539 2540 2541 2542 2543 2544 2545 2546 2547 2548 2549 2550 2551 2552 2553 2554 2555 2556 2557 2558 2559 2560 2561 2562 2563 2564 2565 2566 2567 2568 2569 2570 2571 2572 2573 2574 2575 2576 2577 2578 2579 2580 2581 2582 2583 2584 2585 2586 2587 2588 2589 2590 2591 2592 2593 2594 2595 2596 2597 2598 2599 2600 2601 2602 2603 2604 2605 2606 2607 2608 2609 2610 2611 2612 2613 2614 2615 2616 2617 2618 2619 2620 2621 2622 2623 2624 2625 2626 2627 2628 2629 2630 2631 2632 2633 2634 2635 2636 2637 2638 2639 264

WO 03/100093

PCT/GB03/02307

Method of selecting targets for gene silencing
by RNA interference

Field of the invention

5 The invention relates to the preparation of
double-stranded RNA reagents for use in the specific
down-regulation of gene expression by RNA interference
by hybridisation to scanning arrays of antisense
oligonucleotides.

10

Background to the invention

RNA interference (RNAi) is a process of sequence-
specific gene silencing initiated by double-stranded
RNA (dsRNA) that is homologous in sequence to a region
15 of the silenced gene (Fire, A. Trends Genet. Vol. 15,
358-363, 1999; Sharp, P.A. Genes Dev. Vol. 15, 485-
490, 2001).

Elbashir et al. (Nature, 411, 494-498, 2001) have
20 demonstrated effective RNAi-mediated gene silencing in
mammalian cells using dsRNA fragments of 21
nucleotides in length (also termed small interfering
RNAs or siRNAs). These short siRNAs demonstrate
effective and specific gene silencing, whilst avoiding
25 the interferon-mediated non-specific reduction in gene
expression which has been observed with the use of
dsRNAs greater than 30bp in length (Stark G.R. et al.,
Ann Rev Biochem. 1998, 67: 227-264; Manche, L et al.,
Mol Cell Biol., 1992, 12: 5238-5248). Thus, siRNAs
30 represent promising candidate gene-specific
therapeutic agents, providing an alternative to anti-
sense oligonucleotides.

One potential problem to be overcome in the
35 development of siRNA agents for any given gene is the
selection of an appropriate region of the gene to

WO 03/100093

PCT/GB03/02307

- 2 -

target in order to achieve effective gene-silencing by RNAi.

5 In the field of anti-sense oligonucleotide
technology suitable single-stranded antisense
oligonucleotides for a given gene may be selected with
the use of scanning arrays (Southern, E.M. et al.,
Nucleic Acids Res., 1994, 22(8): 1368-1373; Sohail, M.
and Southern, E.M. "Using oligonucleotide scanning
10 arrays to find effective antisense reagents", *Methods
in Molecular Biology*, vol. 170: *DNA Arrays: Methods
and Protocols*, Ed J.B. Rampal, Humana Press Inc.,
Totowa, NJ; Sohail, M. et al., Nucleic Acids Res.,
2001, 29(10): 2041-2051).

15 The present inventors have now shown that the
sequences selected for use in single-stranded
antisense oligonucleotides on basis of array screening
can also be used as the basis of siRNA reagents that
20 are effective in gene silencing by RNAi.

Description of the invention

In accordance with a first aspect of the
invention there is provided a method of preparing an
25 siRNA reagent for use in gene silencing of a target
gene by RNA interference, which method comprises:

- (a) preparing a scanning array of antisense
oligonucleotides spanning a region of a transcript of
the target gene;
- 30 (b) hybridising to the array labelled transcripts
of the target gene;
- (c) identifying an oligonucleotide within the
array which hybridizes with the labelled transcripts;
and
- 35 (d) preparing an siRNA reagent comprising a
double-stranded RNA of identical sequence to the

WO 03/100093

PCT/GB03/02307

- 3 -

oligonucleotide identified in step (c).

As aforesaid, it is known to use scanning arrays of antisense oligonucleotides in order to identify regions of a particular transcript to target for inhibition of gene expression using antisense oligonucleotides. However, it was surprising to observe that sequences identified using the scanning array as being useful as antisense oligonucleotides may also mediate gene silencing by RNA interference because the mechanisms of action of antisense oligonucleotides and RNA interference are very different (see review by Brantl, S. Antisense-RNA regulation and RNA interference, *Biochem Biophys Acta*. Vol. 1575(1-3), 15-25, 2002).

The term "siRNA reagent" refers to a nucleic acid molecule that is capable of down-regulating expression of a target gene by RNA interference. The characteristics of siRNA reagents are generally known in the art. siRNA reagents generally comprise a region of double-stranded RNA, although as discussed below one or more bases in the double-stranded RNA may be replaced with DNA bases. The double-stranded RNA may be flanked by short single-stranded overhangs, as described below.

Scanning oligonucleotide arrays comprising oligonucleotides complementary to a target region of a mRNA transcript of the gene of interest may be synthesised using the methods which are known in the art and described, for example by Southern E.M. et al., *Nucleic Acids Res.*, 1994, 22(8): 1368-1373 and Sohail, M. and Southern, E.M. "Using oligonucleotide scanning arrays to find effective antisense reagents", *Methods in Molecular Biology*, vol. 170: DNA Arrays:

WO 03/100093

PCT/GB03/02307

- 4 -

Methods and Protocols, Ed J.B. Rampil, Humana Press Inc., Totowa, NJ, the contents of which documents are incorporated herein by reference. A detailed protocol for synthesis of scanning arrays is also given in the accompanying examples.

Scanning arrays are a simple tool that allow combinatorial synthesis of a large number of oligonucleotides on a solid platform (typically glass or polypropylene, see note 1 in accompanying examples) in a spatially addressable fashion, and parallel measurement of the binding of all oligonucleotides complementary to the target mRNA.

The scanning arrays comprise sets of oligonucleotides of various lengths. A series of oligonucleotides, complementary to the target mRNA, is made by sequential coupling of nucleotides to a solid surface. The DNA synthesis reagents are applied to a confined area on the surface of the solid support using a mask (see below). The mask is shifted along the surface after each round of coupling, resulting in a series of oligonucleotides each complementary to a region of the target sequence.

The "gene of interest" may be essentially any gene for which it is desired to develop an siRNA/RNAi reagent. The method of the invention is of general utility, thus the precise nature of this gene (and the mRNAs transcribed therefrom) is not material to the invention. The "target region" may be a sub-fragment of the gene of interest which it is desired to test in order to identify region(s) which may potentially be useful targets for gene silencing by RNA interference. The "target region" may be an arbitrarily chosen sub-fragment of the gene of interest or may have been selected on the basis of an assay for potentially

WO 03/100093

PCT/GB03/02307

- 5 -

suitability as an RNAi target. For example, the "target region" may be one which is relatively accessible in the mRNA transcript because of a relative lack of secondary structure. Regions of mRNA transcripts which are potentially accessible for gene silencing may be identified by RNaseH mapping (see Sohail et al., Nucleic Acids Res., 2001, 29(10): 2041-2051).

10 The scanning arrays will generally containing all complements of the selected target sequence up to a maximum length determined by the size of the template and template displacement used in the synthesis of the scanning array (Southern et al. 1994, *ibid*).

15 Typically, the maximum length of the oligonucleotides in the array will be around 18-20 nt, but this may be varied if required. The short dsRNAs (siRNAs) used for RNA interference are typically 21-23 bp in length, hence it would be appropriate to include similar

20 length sequences in the array.

The scanning arrays are hybridised with a probe which is a transcript of the gene of interest labelled with a revealing label, which may be essentially any

25 type of revealing label which permits visualisation and quantitation of the hybridisation intensity. Radiolabels are particularly preferred. Suitable labelled RNA probes may be conveniently synthesised using standard techniques known in the art (see

30 accompanying examples).

Typically the hybridisation will be carried out at a temperature in the range 20-37°C. Hybridisation at 37°C is particularly preferred, since

35 oligonucleotides which hybridise at this temperature are more likely to be effective *in vivo*. The compositions of standard hybridisation buffers which

WO 03/100093

PCT/GB03/02307

- 6 -

are preferred for use with the arrays are given in the accompanying examples.

Oligonucleotides which hybridise to the array are identified and double-stranded RNA reagents (also referred to herein as RNAi reagents or siRNAs) having identical sequence are synthesised. The inventors have demonstrated by experiment that hybridisation to the scanning array is directly predictive of effectiveness in RNA interference. In this context "identical sequence" is not intended to be interpreted literally as requiring 100% sequence identity. The sequence of the dsRNA may differ slightly from the antisense oligonucleotide (ASO) sequence. For example the length of the dsRNA may be longer or shorter by several nt to optimise performance of the dsRNA. The RNA duplex will preferably be less than 30 bp in length, since duplexes of greater than 30 bp may induce non-specific interferon-mediated effects when introduced into cells *in vivo*. RNA duplexes of 20-27 bp, and typically 20-24 bp, in length are particularly suitable as RNAi reagents. If the dsRNA is made longer than the ASO then the extra sequence may correspond to the "native" sequence of the mRNA. The dsRNA may contain one or more substitute bases in order to optimise performance in RNAi. Substitution of even a single nucleotide may have a profound effect on activity of the RNAi duplex. The dsRNA may further contain non-natural bases or non-natural backbone linkages, for example to enhance stability *in vivo* or enhance resistance to degradation by nucleases. The dsRNA may also include single-stranded overhangs at one or both ends of the duplex. In a particularly preferred embodiment the dsRNA may contain 3' overhanging nucleotides, preferably 3' overhanging thymidines (dTdT) or uridines (UU).

WO 03/100093

PCT/GB03/02307

- 7 -

siRNA reagents may be formed of RNA/DNA chimeras. These chimeras include, for example, the siRNA reagents comprising a double-stranded RNA with 3' overhangs of DNA bases (e.g. dTdT), as discussed
5 above, and also siRNA reagents comprising a double-stranded "RNA" in which one or more of the RNA bases, or even an entire strand, are replaced with DNA bases.

10 In a further embodiment rather than being formed of two separate RNA strands annealed together, the siRNA reagent may comprise a dsRNA having a foldback stem-loop or hairpin structure, wherein the two strands of the dsRNA are covalently linked. RNAs having this structure are typical if the dsRNA is
15 synthesised by expression *in vivo* or by *in vitro* transcription. The precise nature and sequence of the "loop" linking the two RNA strands is generally not material to the invention, except that it should not impair the ability of the double-stranded part of the
20 molecule to mediate RNAi.

The double-stranded RNA will preferably comprise 20-27, or 20-24, consecutive nucleotides of the target mRNA sequence, since duplexes of this length are
25 particularly effective in RNAi.

Double-stranded RNAs may be synthesised *in vitro* using chemical or enzymatic RNA synthesis techniques well known in the art. In one approach the two
30 separate RNA strands may be synthesised separately and then annealed to form double-strands.

In a further embodiment, double-stranded RNAs may be synthesised by intracellular expression from a
35 suitable expression vector. Thus, the invention further provides a method of preparing an expression vector capable of expressing an siRNA reagent for use

WO 03/100093

PCT/GB03/02307

- 8 -

in gene silencing of a target gene by RNA interference, which method comprises:

- (a) preparing a scanning array of antisense oligonucleotides spanning a region of a transcript of the target gene;
- (b) hybridising to the array labelled transcripts of the target gene;
- (c) identifying an oligonucleotide within the array which hybridizes with the labelled transcripts;
- and
- (d) preparing an expression vector capable of expressing an siRNA reagent comprising a double-stranded RNA of identical sequence to the oligonucleotide identified in step (c).

A number of expression vector systems for *in vivo* expression of short double-stranded RNAs for use as RNAi reagents (also referred to as small interfering RNAs, or siRNAs) are known in the art. Generally, siRNAs are expressed as stem-loops, which may be rapidly processed within the cell to produce the "free" siRNA (see review by Tuschl, Nature Biotechnology, Vol. 20(5), 446-448, 2002). Vector systems for expression of siRNAs are often based on RNA Pol III promoters, since these are particularly suited to accurate expression of very short RNA sequences. Suitable vector systems are described in Brummelkamp, T.R. et al., Science, Vol. 296, 550-553, 2002; Lee, N.S. et al., Nature Biotechnology, Vol. 20, 500-505, 2002; Miyagashi, M & Taira, K. Nature Biotechnology, Vol. 20, 497-500, 2002; Paul, C.P. et al., Nature Biotechnology, Vol. 20, 505-508, 2002, the contents of which are incorporated herein by reference.

35

In a further aspect the invention provides a

WO 03/100093

PCT/GB03/02307

- 9 -

method of preparing a pharmaceutical composition comprising an siRNA reagent capable of mediating gene silencing of a target gene by RNA interference, which method comprises:

- 5 preparing an siRNA reagent capable of mediating gene silencing of a target gene by RNA interference according to the method described above and formulating the siRNA reagent into a pharmaceutical composition comprising the siRNA reagent and one or
10 more diluents, excipients or carriers.

- siRNA reagents may be formulated into pharmaceutical compositions comprising a
15 therapeutically effective amount of the siRNA nucleic acid in combination with any standard physiologically and/or pharmaceutically acceptable carriers known in the art. "Pharmaceutically acceptable" means a non-toxic material which does not interfere with the activity of the pharmaceutically active ingredients in
20 the composition. "Physiologically acceptable" refers to a non-toxic material that is compatible with a biological system such as a cell, tissue or organism. Physiologically and pharmaceutically acceptable carriers may include diluents, fillers, salts,
25 buffers, stabilizers, solubilizers etc.

- For delivery into cells in vivo siRNAs may be formulated with lipid-based carriers including, for example, oil-in water emulsions, micelles, and
30 liposomes. Liposomes are the most preferred carriers, and there use is well known in the art. Liposomes are commercially available from Gibco BRL, for example, as LIPOFECTIN™ and OLIGOFECTAMINE™, which are formed of cationic lipids. Methods for making liposomes are
35 well known in the art and have been described in many publications. Liposomes may be targeted to a particular tissue by coupling the liposome to a

WO 03/100093

PCT/GB03/02307

- 10 -

particular tissue by coupling the liposome to a specific ligand, such as a monoclonal antibody, sugar, glycolipid or protein.

5 Liposomes may also be used to deliver vectors encoding siRNAs. In the field of human gene therapy it is well known to deliver expression vectors, such as plasmids, via nucleic acid-liposome complexes.

10 For use in human therapy, pharmaceutical compositions including the siRNAs of the invention will be administered to a patient in need of treatment in a "therapeutically acceptable amount". A therapeutically acceptable amount is an amount of a
15 pharmaceutical preparation that alone, or together with further doses, produces the desired response in the condition being treated. The precise amount of the composition administered will, however, generally be determined by a medical practitioner, based on the
20 circumstances pertaining to the disorder to be treated, such as the severity of the symptoms, the composition to be administered, the age, weight, and response of the individual patient and the chosen route of administration.

25 The invention will be further understood with reference to the following experimental protocols, together with the accompanying Figures, in which:

30 Figure 1 shows representative plots of hybridisation intensity across the array for (a) 15mers and (b) 18mers probed with labelled IGF1R mRNA.

35 Figure 2 illustrates the effect of ASOs and RNAi on IGF1R levels in MDA-231 human breast cancer cells. The cells were transfected at 30-40% confluence with phosphorothioate ASOs complexed with the lipid

WO 03/100093

PCT/GB03/02307

- 11 -

Cytofectin (Glen GSV). After 48hr the cells were lysed and equivalent amounts of soluble protein were separated by SDS-PAGE and immunoblotted for IGF1R and β -tubulin (loading control). The intensity of the autoradiographic bands was quantified by densitometry, and IGF1R levels were corrected for loading differences. The results are shown as % IGF1R level of that in cells transfected with the same concentration of an appropriate control. This was a scrambled control oligonucleotide for ASOs, and an inverted RNA duplex for RNAi.

Figure 3 illustrates the effect of ASOs and RNAi on IGF1R and IR levels in MDA-231 human breast cancer cells and ME melanoma cells. Cells were transfected using either Cytofectin (C) or Oligofectamine (O; Gibco BRL). After 48hr the cells were lysed and IGF1R and IR levels were determined by immunoblotting.

a) MDA-231 breast cancer cells transfected with ASO 2 (or scrambled control, Scr) or RNAi or inverted RNA duplex control (InvRNA) at 20 or 200nM.

b) MDA-231 cells were transfected with Oligofectamine and RNAi or Inverted control (InvRNA) at 0.1 - 10nM.

c) ME melanoma cells were transfected with Oligofectamine and RNAi or Inv control duplex at 5 - 500 nM.

Figure 4 illustrates the effects of various RNAi duplexes in MDA-231 breast cancer cells. MDA-231 cells were transfected with oligofectamine and 21mer RNA duplexes at 0.5, 5 and 50nm. After 48hr IGF1R expression was analysed by immunoblotting (panel (a)). Panel (b) is a graphical illustration, results are presented as % IGF1R level of that in cells transfected with the same concentration of an inverted control RNAi.

WO 03/100093

PCT/GB03/02307

- 12 -

Figure 5 illustrates the activity of RNAi duplexes of 18, 21 or 24 nt in MDA-231 breast cancer cells. Panel (a) is an immunoblot showing the effect of various RNAis on IGF1R expression at 0.5 and 5nM; panel (b) is a graphical illustration showing the effect of RNAis of varying length, results are presented as % IGF1R level of that in cells transfected with the same concentration of an Inv2 control RNAi.

Figure 6 illustrates the effect of RNAi duplexes on Akt phosphorylation in ME melanoma cells. ME cells were transfected with 100nM RNA duplexes. After 48hr the monolayers were disaggregated and half of each culture was treated with 10nM IGF-1 for 30 min. The cells were lysed and lysates were analysed by immunoblotting for IGF1R, phospho-Ser473-Akt and total Akt.

Figure 7 illustrates growth in vivo in C57BL mice of B16 melanoma cells transfected with RNA duplexes. B16 melanoma cells were transfected with 200nM duplexes (RNA22=R2,22Inv=Inv2) or oligofectamine alone (OF) for three consecutive days. On the 4th day cells were injected into the flanks of C57BL mice, using 2.5×10^5 cells/mouse, using groups of 5 mice. Tumours were measured in 2 dimensions every 2-3 days. Tumour volumes were calculated as $n(a \times b^2)/6$ where a is the larger and b the smaller dimension.

Figure 8 shows the complete cDNA sequence for human IGF1R. The region evaluated using the scanning array is underlined.

Example 1-Synthesis of a scanning array

In the following protocol references to Figures and Tables from "Sohail and Southern" refer to Sohail,

WO 03/100093

PCT/GB03/02307

- 13 -

M. and Southern, E.M. "Using oligonucleotide scanning arrays to find effective antisense reagents", *Methods in Molecular Biology*, vol. 170: DNA Arrays: Methods and Protocols, Ed J.B. Rampil, Humana Press Inc., Totowa, NJ.

Using a diamond-shaped or a circular reaction mask (Sohail and Southern, Fig. 1), it is possible to create arrays comprising sets of oligonucleotides of all lengths from monomers up to a maximum in a single series of couplings. The maximum length of oligonucleotides synthesised depends upon the ratio of the diagonal (for a diamond-shaped mask) or diameter (for a circular mask) of the mask to the displacement at each coupling step. For example, a diamond-shaped mask of 40 mm diagonal will produce 10-mers, 16-mers, or 20-mers using step sizes of 4 mm, 2.5 mm, or 2 mm, respectively. A diamond-shaped template creates a series of small diamond-shaped cells. The longest oligonucleotides are found along the centre line and the monomers are located at the edge (Sohail and Southern, Fig. 1). A circular template creates cells that differ in shape: along the centre line, they are lenticular, but off this line, they form a four-cornered "spearhead" that diminishes in size towards the edge. The arrays as synthesised are symmetrical above and below the centre line of the template and each oligonucleotide is represented twice allowing for duplicate hybridisation measurements.

For each length of oligonucleotides s , there are $N-s+1$ s -mers covering a total length of N bases. For example, if a 150 nt long sequence is covered in a 150 step synthesis, there will be 150 monomers and 131 20-mers. The last 20 positions in the sequence will be represented by shorter oligonucleotides only; in this

WO 03/100093

PCT/GB03/02307

- 14 -

case, from 19-mer to monomer. Therefore, for making 200 20-mers, an additional 19 nt synthesis steps need to be added at the end, i. e., total coupling steps = $N_s - 1$.

5

Materials

Derivatisation of Glass

1. Glass cylinder and apparatus shown in Sohail and Southern, Fig. 3.
- 10 2. Glass sheets of required dimension (3 mm thick: Pilkington, UK).
3. 3-Glycidioxypropyl trimethoxysilane (98% v/v: Aldrich).
4. Di-isopropylethylamine (99.5 % v/v: Aldrich).
- 15 5. Xylene (AnalaR: Merck).
6. Hexaethylene glycol (97 % v/v: Aldrich).
7. Sulfuric acid (AnalaR: Merck).
8. Ethanol (AnalaR: Merck).
9. Ether (AnalaR: Merck).
- 20 10. Water bath at 80°C.

Making Reaction Masks

1. Stainless steel or aluminium square metal piece or PTFE (Teflon). Dimensions of the workpiece may vary
- 25 according to the size of the mask.
2. A centre lathe or a horizontal milling machine.
3. A drilling machine.
4. Abrasive paper from ~ P600 to P1200 (3M Inc., USA) and polishing grade crocus paper (J. G. Naylor & Co.
- 30 Ltd., Woodley, Stockport, Manchester, England).

Making Scanning Arrays

1. Solid support (derivatised glass or aminated polypropylene (Beckman Coulter, Inc., USA)).
- 35 2. DNA synthesiser (ABI).
3. A reaction mask of desired shape and size and assembly frame (see Sohail and Southern Fig. 4 for the

WO 03/100093

PCT/GB03/02307

- 15 -

assembly).

4. DNA synthesis reagents: standard dA, dG, dC and T phosphoramidites, oxidizing agent, acetonitrile, activator solution, deblock solution (all from Cruachem).

5. Reverse phosphoramidites bought from Glen Research

Deprotection of Arrays

1. Assembly for constructing deprotection bomb as shown in Sohail and Southern Fig. 5. The assembly consists of a high density polyethylene (HDPE) chamber, 4 mm thick silicon rubber gasket and a stainless steel plate of the dimensions of the HDPE chamber, and stainless steel M8 nuts and bolts.

2. 30% ammonia solution (AnalaR: Merck).

3. Water bath at 55°C.

In Vitro Transcription

1. Template DNA (at ~1 mg/mL).

2. T7 or SP6 RNA polymerase, transcription buffer (5X transcription buffer is, 200 mM Tris-HCl pH 7.9, 30 mM MgCl₂, 10 mM spermidine, 50 mM NaCl), recombinant RNasin®, 100 mM DTT and nuclease-free distilled water (Promega).

3. [α -³²P] UTP (3000 Ci/mmol) or [α -³³P] UTP (2500 Ci/mmol) (Amersham).

4. rNTPs (Pharmacia): ATP, GTP, CTP stored as 10 mM solution, and UTP as 250 mM solution in nuclease-free distilled water. Store all reagents at -20°C.

Quantitation of Transcripts

1. Scintillation counter (e. g., Beckman LS 1710).

2. Scintillation vials and scintillation fluid (Amersham).

Hybridisation, Imaging and Analysis

1. Hybridisation buffer (1M NaCl, 10 mM Tris-HCl pH

WO 03/100093

PCT/GB03/02307

- 16 -

- 7.4, 1 mM EDTA, 0.01 % SDS (w/v)) (see Note. 2).
2. 50-100 fmol radiolabelled transcript.
3. A glass plate of the size of the array when using
an array made on glass, a moist chamber (a large
plastic or glass lidded box containing wetted paper
towels) and an incubator set to the desired
temperature.
4. A hybridisation tube and oven used in standard
Southern hybridisation (e. g., Techne) when using an
array made on polypropylene.
5. Esco rubber tubing of OD 1 mm (Sterlin) for use in
Section 3.8.
6. Storage phosphor screen (Fuji or Kodak).
7. PhosphorImager or STORM (Molecular Dynamics).
8. A SUN Solaris work station for image analysis and
the computer software xvseq (L. Wand and J. K. Elder,
unpublished) (available by anonymous ftp at
ftp://bioch.ox.ac.uk/pub/xvseq.tar.gz).

Stripping of Arrays

1. Stripping solution (100 mM sodium
carbonate/bicarbonate buffer pH 10, 0.01 % SDS (w/v))
(see Note 3).
2. Geiger-Müller counter (Mini-Instruments Ltd.).

Methods

Derivatisation of Glass

1. Prepare a mixture of di-isopropylethylamine,
glycidoxypopyl trimethoxysilane and xylene (1: 17.8:
69, v/v/v) in a glass cylinder and completely immerse
the glass plates in the mixture. Incubate as shown in
Sohail and Southern Fig. 3 at 80°C for 9 h.
2. Remove the plates, allow them to cool to room
temperature, and wash with ethanol and then with ether
by squirting the liquid from a wash bottle.
3. Incubate the plates in hexaethylene glycol
containing a catalytic amount of sulfuric acid (~ 25

WO 03/100093

PCT/GB03/02307

- 17 -

mL/L) at 80°C for 10 h, with stirring.

4. Remove the plates, allow them to cool to room temperature, and wash with ethanol and ether. Air dry and store at -20°C.

5

Machining of Masks

1. Both stainless steel or aluminium can be used to make diamond-shaped and circular reaction masks. Circular masks are made using a centre lathe and diamond-shaped masks using a horizontal milling machine (see Note 4).
2. To make a diamond shaped mask from metal, hold the work piece at an angle of 45° to the axis of the bed of the milling machine (the diagonal of the diamond running parallel to the axis of the bed).
3. Machine the cavity to the required depth (generally between 0.5-0.75 mm) to create a reaction chamber. Machine the outer lands to a depth of approximately 0.5 mm to form the sealing edge (0.3-0.5 mm wide) (Sohail and Southern Fig. 4).
4. Using the smallest possible diameter cutter (~1.5 mm), radius the internal corners of the reaction chamber.
5. Finish the sealing edge by polishing flat with successively finer grades of wetted abrasive paper (from ~ P600 to P1200) and finally with a polishing grade crocus paper.
6. Drill holes of 1.08 mm diameter for reagent inlet and outlet, respectively, at the bottom and the top of the reaction chamber (in the corners of the diamonds). Inlet and outlet connections to the DNA synthesiser are made using standard 19SWG syringe needles (1.1 mm diameter) with chamfered ends ground off and de-burred (see Note 5).

35

Fabrication of Arrays

1. Cut glass or polypropylene to the correct size.

WO 03/100093

PCT/GB03/02307

- 18 -

- The process of making an array is the same when using either glass or polypropylene. Polypropylene has to be mounted on a glass plate, e. g., 3 mm thick soda glass (see Note 6). The total area covered by an array for N bases using a mask of diagonal or diameter D mm and step size l mm is $N \times l + D$ mm. 2-3 mm are added to margins to allow easy manipulations.
2. Fix the assembly (Sohail and Southern Fig. 4) to the front of a DNA synthesiser and connect its inlet and outlet to the synthesiser's reagent supply.
3. Program the DNA synthesiser with an appropriate synthesis cycle. A slightly modified cycle is used, for example, the one given in Table 1 (Sohail and Southern). Also check all the reagent bottles.
4. Enter the sequence (antisense strand) in 5' to 3' direction.
5. Mark the first footprint of the reaction mask on the support by placing it against the mask on the assembly in the desired starting position (see Sohail and Southern Fig. 6). A knife is used to make notches in polypropylene: A diamond scribe can be used to mark glass.
6. Tighten the plate against the mask with the pressure clamp to produce a seal (Sohail and Southern Fig. 4b). Sufficient pressure is applied to stop leakage (~ 500-800 Newton force) but not enough to create indentations in the polypropylene surface which can lead to leakage of reagent from the mask during subsequent synthesis steps.
7. Start the DNA synthesiser to go through the pre-programmed cycle to couple the appropriate nucleotide. The first condensation on the substrate is of base at the 3' end of the sequence.
8. After completion of the step during the interrupt (see Sohail and Southern Table 1 and Note 7), slacken the pressure clamp and move the plate one increment (Sohail and Southern Fig. 4b).

WO 03/100093

PCT/GB03/02307

- 19 -

9. Tighten the pressure clamp and start the synthesiser for the next nucleotide in the sequence. Continue the process until the full sequence length is synthesised (see Note 8).

5

Deprotection of Arrays

1. Place the glass or polypropylene array(s) into the HDPE chamber (see Note 9) and add 30% ammonia into the chamber to cover the array(s) completely.
- 10 2. Place the silicon rubber gasket around the rim of the chamber and the stainless steel plate on top of the gasket.
3. Place bolts through the metal plate, the gasket and the HDPE chamber, and tighten.
- 15 4. Incubate in a water bath at 55°C for 12-18 h in a fume hood.
5. Cool the assembly to 4°C before opening. The arrays are ready to be used in hybridisation at this stage.

20

Preparing and Quantifying Radiolabelled Transcripts

1. Set an *in vitro* transcription reaction (20 µL) by adding the following components to a microfuge tube at room temperature.
 - 25 5 X transcription buffer 4 µL
 - 100 mM DTT 2 µL
 - RNAasin™ 20 U
 - 10 mM ATP, GTP, and CTP 1 µL each
 - 250 mM UTP 1 µL
 - 30 Template DNA 2-3 µL (see Note 10)
 - [α-32P]UTP or [α-33P]UTP 2 µL
 - T7 or SP6 RNA polymerase 20 U
 - Total Volume 20 µL
 - Mix and incubate at 37°C for 1 h.
- 35 2. Remove 1 µL for quantitation (see below).
3. Remove unincorporated label by Sephadex® G25 column chromatography (see Note 11).

WO 03/100093

PCT/GB03/02307

- 20 -

4. Save 1 μL of the purified transcript for quantitation (see below).
5. Check the integrity of the transcript by denaturing polyacrylamide gel electrophoresis (10).
- 5 6. Add 10 μL of the scintillation fluid to the samples saved in Steps 2 and 4.
7. Mix by vortexing and count the samples in a scintillation counter for 1 min.
8. Calculate the per cent incorporation:
10 % incorporation = $\frac{\text{incorporated cpm} \times 100}{\text{total cpm}}$
9. Calculate the amount of RNA made:
Amount of $[\alpha\text{-}^{32}\text{P}]\text{UTP}$ = $\frac{20 \mu\text{Ci}}{3000 \mu\text{Ci/nmol}}$ = 6.6×10^{-3} nmol
- 15 Amount of cold UTP = $1 \mu\text{L} \times 250 \mu\text{M} = 0.250$ nmol
Total UTP = $6.6 \times 10^{-3} + 0.25 = 0.256$ nmol
For a reaction with 50 % incorporation, the amount of UTP incorporated
= $0.256 \text{ nmol} \div 2 = 0.128$ nmol
- 20 Supposing equal incorporation of all four nucleotides, total nucleotides incorporated
= $0.128 \text{ nmol} \times 4 = 0.512$ nmol
Amount of full-length transcript
= $0.512 \div \text{total length of transcript}$
- 25 Hybridisation to Arrays Made on Polypropylene
1. Place the array in the hybridisation tube, coiling it in a spiral.
2. Dilute the radiolabelled transcript in an
30 appropriate volume (10-20 mL depending upon the size of the array and the hybridisation tube) of hybridisation buffer. The mix should cover the array along the length of the tube.
3. Place items 1 and 2 in the oven at desired
35 temperature for 30 min. Also put approximately 100 mL of the hybridisation buffer in the oven: this is to be used to wash the array at the end of hybridisation.

WO 03/100093

PCT/GB03/02307

- 21 -

4. Pour the hybridisation mix into the tube containing the array and hybridise for 3-4 h.
 5. Remove the hybridisation mix. Briefly wash the array with the hybridisation buffer from Step 3, air dry, cover with cling film and expose to a storage phosphor screen for 16-20 h (see Note 12).
 6. Scan the screen on PhosphorImager or STORM and analyse the image using xvseq (see below).
-
- 10 Hybridisation to Arrays Made on Glass
 1. Clean the non-array glass plate with acetone and ethanol to ensure it is grease-free and siliconise it by treatment with dimethyl dichlorosilane solution and place it in lidded box. Also place moist paper towel in the box.
 2. Dilute the radiolabelled transcript in an appropriate volume of the hybridisation buffer (for example, for an array of 250 mm x 50 mm use 500-750 μ L).
 - 20 3. Place items 1 and 2 in an incubator at desired temperature for 30 min. Also put approximately 100 mL of the hybridisation buffer in the oven: this is to be used to wash the array at the end of hybridisation.
 4. Using a micropipette, pipette the hybridisation mix in a line evenly along the length of the non-array glass plate, avoiding formation of air bubbles.
 - 25 5. Starting at one end, carefully place the scanning array (face down) on top of the hybridisation mix. The mix will spread out and form a thin film between the two plates. Incubate for 3-4 h.
 - 30 6. Separate the plates from each other and wash the array plate with hybridisation buffer to remove unbound mix. Drain the plate, air dry, cover with cling film and expose to a storage phosphor screen for 16-20 h.
 - 35 7. Scan the screen on PhosphorImager or STORM and analyse the image using xvseq (see below).

WO 03/100093

PCT/GB03/02307

- 22 -

Alternative Hybridisation Protocol for Glass or Polypropylene

1. Assemble with clips the array plate (or polypropylene array pasted with PhotoMount™ on a glass plate) and the non-array plate, using rubber tubing as spacers on two sides.
2. Dilute the radiolabelled transcript in approximately 5-10 mL of hybridisation buffer.
3. Follow Step 3, Protocol 3.7.
4. Inject the hybridisation mix into the space between the two plates with a needle and syringe.
5. Incubate the assembly in horizontal position at desired temperature.
6. Follow Steps 6 and 7, Protocol 3.7.

Image Analysis

The hybridisation images are analysed using xvseq (see Sohail and Southern Fig. 7). This program reads and displays images generated by a PhosphorImager or STORM and can also perform standard image manipulation such as scaling, clipping and rotation. Although visual inspection of an image reveals the results generally, computer-aided analysis is needed to obtain quantitative information about hybridisation intensities and the oligonucleotide sequences that generated them. xvseq calculates and displays integrated intensities of the array oligonucleotides, each of which corresponds to an image cell formed by intersection of overlapping array templates.

The user can specify the template size, shape and location, step size between successive templates, as well as the sequence that was used to make the array. The template grid is superimposed on the image and the template parameters are adjusted interactively to achieve correct and accurate registration of the grid with the hybridisation pattern. It can be difficult to

WO 03/100093

PCT/G803/02307

- 23 -

achieve precise registration by reference to the hybridisation pattern alone, especially, if the signals at either edge of the array are weak or undetectable. Avoid placing the template grid so that it appears to be registered but is in fact misaligned by one or more template steps. Registration can be aided by the use of fixed reference points on an array such as those shown in Sohail and Southern Fig. 6.

10 Stripping of Arrays

1. The arrays can be used several times. To strip, heat an appropriate volume of the stripping solution to 90°C in a glass beaker.
2. Immerse the array in the hot stripping solution and stir for 1-2 min.
3. Remove the array and monitor with a Geiger counter to confirm that most of the radiolabel has been removed. Repeat steps 1-2 if radioactivity on the surface of the array is still detectable.
4. Allow the array to cool down to room temperature and wash it thoroughly with nuclease-free distilled water, 70 % (v/v) ethanol and finally with absolute ethanol.
5. Air dry and store the array at -20°C until future use.

Notes

1. The choice of array substrate material and attachment chemistry is important for making high quality arrays. A flat, impermeable surface is required for *in situ* synthesis of arrays. Glass has a number of favourable qualities, including its wide availability, smooth surface, transparency, chemical stability and compatibility with the use of both radiolabelled or fluorescence labelled nucleic acids targets. Glass is chemically derivatised as described in the methods section to produce a hexaethylene

WO 03/100093

PCT/GB03/02307

- 24 -

glycol linker which has a terminal -OH group that allows condensation of nucleotide phosphoramidites (Maskos, U., and Southern, E. M. (1992) Nucleic Acids Res. 20, 1679-1684.). Similarly, polypropylene also has favourable physical and chemical properties. Polypropylene is aminated to produce amine groups (Matson, R. S., Rampal, J. B., and Coassin, P. J. (1994) Anal. Biochem. 217, 306-310.) that also allow synthesis to oligonucleotides using standard CE nucleotide phosphoramidites.

For the array fabrication method described here, it is important that a tight seal is formed between the substrate material and the reaction mask. Metals form tight seal with polypropylene but not with glass. PTFE seals well against both glass and polypropylene.

2. 1M NaCl is used routinely. Alternative buffers are: (i) 1 M NaCl, 5-10 mM MgCl₂, 10 mM Tris-HCl pH 7.4, 1 mM EDTA, 0.01 % SDS (w/v), and (ii) 150 mM NaCl, 10 mM MgCl₂, 10 mM Tris-HCl pH 7.4, 1 mM EDTA, 0.01 % SDS (w/v).

3. Addition of more than 0.01%SDS can damage arrays.

4. Circular masks can also be made from PTFE (Teflon). Diamond-shaped masks are more difficult to make with PTFE by the machining process but can be made by pressure moulding in a hydraulic press (~ 150 ton force) using a pre-machined die.

5. Holes should be made as close as reasonably possible to the sealing edge without damaging it. Care must be taken to de-burr fully the holes at the point of entry into the reaction chamber. For PTFE masks the holes should be 1.0 mm diameter which make virtually 100% leak-tight seal. In the case of metal masks, the 0.02 mm interface indicated above also provides a

WO 03/100093

PCT/GB03/02307

- 25 -

leak-tight seal without the use of any additional sealer. Care must be taken not to insert the end of the syringe needle into the reaction chamber void.

5 6. Unlike glass, polypropylene is not rigid and thus
needs to be mounted on a solid, flat surface for its
precise movement against the reaction mask during
synthesis. Even mounting of polypropylene on glass is
important to produce a good seal between the sealing
10 edge of the reaction mask and the polypropylene
surface. Glass used must be clean and free from dust
particles because they can cause bulging of the
polypropylene which can hinder the formation of a
proper seal. A very thin layer of PhotoMount™ (3M
15 Inc, USA) which can be used to paste polypropylene to
glass, should be sprayed on glass and not
polypropylene.

20 7. At the start of each synthesis cycle, an
interrupt step can be introduced to halt the process
at the first step of the next nucleotide condensation
cycle to allow the operator to move the plate and
restart the program. Alternatively, a long wait step
at the beginning of the program can be introduced (see
25 Sohail and Southern Table 1) if the operator does not
wish to use the interrupt step. The operator is also
advised to consult the user's manual for the DNA
synthesiser.

30 8. With the use of standard phosphoramidites in the
synthesis, the oligonucleotides are attached to the
solid support at their 3' ends. Reverse
phosphoramidites can be used to make oligonucleotides
that are attached at their 5' ends.

35

Iodine is used as an oxidising agent. At lower
temperatures it will take longer to reach the top of

WO 03/100093

PCT/GB03/02307

- 26 -

the reaction cell. Iodine can also be replaced with sulfurising agent (Cruachem) to make arrays of phosphorothioate oligonucleotides.

- 5 9. When using the standard phosphoramidites, the exocyclic amines of the bases are protected chemically to prevent side reactions during synthesis. These protecting groups need to be removed from the coupled bases before hybridisation.

10

Before deprotection, detach the polypropylene arrays from glass by peeling from one end. PhotoMount can be removed with ethanol, acetone or dichloromethane.

15

10. An internally radiolabelled RNA is used as target to hybridise to a scanning array which is generated by in vitro transcription, carried out in the presence of [α -³²P]UTP or [α -³³P]UTP (or [α -³²P]CTP) using an appropriate DNA template. A plasmid containing the desired DNA fragment under the transcriptional control of a T7 or SP6 promoter (such as pGEM: Promega) can be used as template. The plasmid is linearised with an appropriate restriction endonuclease to produce transcripts of defined length without contaminating vector sequence. Alternatively, a template with T7 or SP6 RNA promoter can also be generated using the polymerase chain reaction: primers are used to amplify the required fragment from a plasmid, genomic DNA or cDNA, such that the sense primer has a T7 or SP6 promoter leader sequence (Sohail and Southern Table 2) added at the 5' end.

25

30

11. Sephadex® G25 columns are available from several commercial suppliers including Promega and Pharmacia. Spin columns made in-house, as described in (Sambrook, J., Fritsch, E. F., and Maniatis, T. (1989) Molecular

35

WO 03/100093

PCT/GB03/02307

- 27 -

Cloning: A Laboratory Manual. Cold Spring Harbor, NY.), can also be used.

12. For hybridisation below 37°C, care must be taken not to touch the plates because this can lead to melting of short duplexes. For hybridisation below room temperature, the cling film and the phosphor screen must be cooled to hybridisation temperature and exposed at the same temperature.

10

Example 2-Use of scanning array to select RNAi reagents

- The following is give by way of example of the utility of scanning arrays in the selection of RNAi reagents. The invention is of general utility in the selection of RNAi reagents, thus it is not in any way intended to limit the scope of the invention to this specific example.

- Scanning arrays complementary to the region of the IGF1R mRNA from position 537-685 were prepared using the standard techniques described above. The maximum length of the oligonucleotides in the array was either 18 or 20 nt.

25

- In this study hybridisation to the arrays was carried out at physiological temperature (37°C) in addition to room temperature (23°C), in order to select sequences which are more likely to have activity in intact cells. The arrays were probed with labelled IGF1R mRNA, and also with labelled insulin receptor mRNA in order to identify oligonucleotides which have high specificity for IGF1R mRNA.

- IGF1R template/probe preparation

Human IGF1R cDNA in plasmid pCVN was a generous gift from Renato Baserga. The 5' region of IGF1R cDNA

WO 03/100093

PCT/GB03/02307

- 28 -

was cloned into vector pBluescript KS- (Stratagene) using restriction sites HindIII (pCVN-derived site at 5' end of IGF1R cDNA) and Asp718 (cuts IGF1R cDNA at position 1581). This construct (template 1, 1.6kb) included approximately 100bp of polylinker sequence between the T7 promoter and the start of the IGF1R sequence. It was thought that this extraneous vector-derived sequence might influence folding of the transcript. Therefore this region was shortened by restriction digest using NotI, filling in with Klenow enzyme, digestion with EcoRV and re-ligating to make construct 2 (1.6kb Δ 55). Both constructs were linearised at the 3' end of the IGF1R insert using Asp718, to make templates for in vitro transcription. End-labelled transcripts 1 (1.6kb) and 2 (1.6kb Δ 55) (see Figure 1) were generated using T7 RNA polymerase and γ -³²P-GTP (Amersham Pharmacia).

Results

Representative plots of hybridisation intensity across the array for 15mers and 18mers probed with labelled IGF1R mRNA are shown in Figure 1. The plots are annotated to show the position of selected individual oligonucleotide sequences. In each case the sequence given corresponds to the sense strand of the IGF1R mRNA.

Relative hybridisation intensities for selected oligonucleotides were calculated relative to the most strongly hybridising sequence.

WO 03/100093

PCT/GB03/02307

- 29 -

Table 1: Relative hybridisation intensity

Oligo	Sequence	Hyb intensity	Rel hyb intensity
ASO2	ggctcttcacacatcgg	56,026	1.00
ASO4	gcggtagttgtactcattgt	>33,162	>0.59
ASO1	ggcttctcctccatgggc	15583	0.28
ASO3	attgttgatgggtgttcttct	>7048	>0.13
ASO5	gtgggtccagcagcggtagtt	>4150	>0.07
ASO6	cttattccccacaatgta	560	0.01

Example 2-Synthesis of RNAi duplexes

Double-stranded RNA duplexes for RNAi corresponding in sequence to certain of the ASOs identified on the basis of the array screening as hybridising strongly or weakly with IGF1R mRNA were synthesised, together with corresponding inverted controls, mouse homologs and mutant variants (see Table 2). RNA oligonucleotides were synthesised and HPLC purified at Cruachem, Glasgow. Lyophilised oligoribonucleotides were reconstituted in nuclease-free water and diluted to 50µM. Complementary strands were annealed in 100mM potassium acetate, 30mM Hepes-KOH pH 7.4, 2mM magnesium acetate, as described (Elbashir et al., Nature, 2001, www.dharmacon.com) to give a final concentration of 20µM duplex. Duplex formation was checked by electrophoresis through 5% low-melting point agarose (NuSieve GTG, BioWhittaker Molecular Applications) in 1xTBE.

WO 03/100093

PCT/GB03/02307

- 30 -

Table 2: RNAi duplex sequences

5	R2	GCCGAUGUGUGAGAAGACCTT TTCGGCUACACACUCUUCUGG	SEQ ID NO:23
	R2 (18mer)	GCCGAUGUGUGAGAAGTT TTCGGCUACACACUCUUC	SEQ ID NO:24
	R2 (24mer)	GCCGAUGUGUGAGAAGACCACCTT TTCGGCUACACACUCUUCUGGUGG	SEQ ID NO:25
10	R2 (27mer)	GCCGAUGUGUGAGAAGACCACCAUCTT TTCGGCUACACACUCUUCUGGUGGUAG	SEQ ID NO:26
	Inv2	CCAGAAGAGUGUGUAGCCGTT TTGGUCUUCUCACACAUCGGC	SEQ ID NO:27
	Mut2	GCCGAUGUGUGUGAAGACCTT TTCGGCUACACACUCUUCUGG	SEQ ID NO:28
15	MouseR2	GCCCAUGUGUGAGAAGACCTT TTCGGGUACACACUCUUCUGG	SEQ ID NO:29
	R6	CUACAUUGUGGGGAUAAGTT TTGAUGUAACACCCCUUAUUC	SEQ ID NO:30
	Inv6	GAAUAAGGGGUGUACAUCTT TTCUUAUCCCCACAAUGUAG	SEQ ID NO:31
	MouseR6	CUACAUUGUGGGGAACAAGTT TTGAUGUAACACCCCUUGUUC	SEQ ID NO:32

For comparison purposes the following antisense
oligonucleotides were also synthesised:

20

WO 03/100093

PCT/GB03/02307

- 31 -

Table 3: Antisense oligonucleotides

	ASO1 (18mer)	GGCTTCTCCTCCATGGTC	SEQ ID NO:1
	ASO1 (20mer)	CGGCTTCTCCTCCATGGTCC	SEQ ID NO:2
5	Mouse ASO1	GGGCTTCTCCTCCAATGTCC	SEQ ID NO:3
	Scr1	TCTCCCGGACTTGCTCCGC	SEQ ID NO:4
	Mouse Scr1	CTGTTACCGTGCACCCTGT	SEQ ID NO:5
	ASO2 (18mer)	GGTCTTCTCACACATCGG	SEQ ID NO:6
	ASO2 (20mer)	TGGTCTTCTCACACATCGGC	SEQ ID NO:7
10	Mouse ASO2	TGGTCTTCTCACACATGGGC	SEQ ID NO:8
	Scr2	CCTGTACGCGTTGATCTCCA	SEQ ID NO:9
	Mouse Scr2	AGTCGCCTAGTCGAGTCCTT	SEQ ID NO:10
	ASO3	ATTGTTGATGGTGGTCTTCT	SEQ ID NO:11
	Scr3	TCATGCTTGTTGATGTGGTT	SEQ ID NO:12
15	ASO4	GCGGTAGTTGTACTCATTGT	SEQ ID NO:13
	Scr4	TGTCGTTGAGTTGATCGTA	SEQ ID NO:14
	ASO5	GTGGTCCAGCAGCGGTAGTT	SEQ ID NO:15
	Scr5	GACGTTAGCGTGCGATGTGC	SEQ ID NO:16
	ASO6	CTTATTCCCCACAATGTA	SEQ ID NO:17
20	Mouse ASO6	CTTGTTCCTCCCCACAATGTA	SEQ ID NO:18
	Scr6	CATAGCACATTCTCTCA	SEQ ID NO:19
	Mouse Scr6	ACCGTTCATCAGTTCACT	SEQ ID NO:20
	TSS	TCCTCCGGAGCCAGACTT	SEQ ID NO:21
25	ScrTSS	CAGCTACTCGCATGCTGC	SEQ ID NO:22

Example 3-Testing of RNAi duplexesCell culture and transfection

30 The human and murine cell lines used in this study were cultured in RPMI 1640 plus 10% FCS at 37°C in a humidified atmosphere of 5% CO₂. The cultures

WO 03/100093

PCT/GB03/02307

- 32 -

were negative when tested for *Mycoplasma* infection. Cultures were passaged the day before transfection to achieve 30-50% confluence the following day. Initial transfections of ASOs used 1-2.5 $\mu\text{g/ml}$ Cytofectin (Glen Research, Sterling, VA), according to the manufacturer's instructions. Latterly Oligofectamine (Life Sciences) has been used for all ASO and RNAi transfections. The method was as described (see manufacturer's instructions and www.dharmacon.com) with minor modifications. For transfection of monolayers seeded in 6 well plates, 5 μl phosphorothioate oligonucleotide or RNA duplex at 100x final concentration were mixed with 250 μl serum-free Optimem (Gibco-BRL). In a separate tube 2 μl Oligofectamine was mixed with 68 μl Optimem. After 10min incubation at room temperature the contents of the two tubes were mixed and incubated for a further 25min at room temperature. Monolayers were washed with 1-2ml Optimem. To the cells were added 175 μl Optimem followed by the 325 μl complexes. Volumes were scaled up by factors of 2.22 or 6.05 for transfection in 60mm or 100mm dishes respectively. Cultures were incubated at 37°C for 4hr, and then 50 μl FCS and 3.5ml RPMI plus 5% FCS were added to each well. After 48hr incubation effects on IGF1R and IR expression were determined by immunoblotting. Some transfected cultures were disaggregated using 3mM EDTA in PBS for 2-4min, washed in serum-free RPMI, divided into two aliquots and treated with 10nM Long R3-IGF-I (GroPep, Adelaide) or diluent for 30min at 37°C prior to lysis.

Immunoblotting

IGF1R expression was assessed by immunoblotting as previously described (Macaulay et al 2001). After washing in ice-cold PBS, cells were lysed in 50mM Hepes pH 7.5, 100mM NaCl, 10mM EDTA, 1% Triton-X-100, 4mM sodium pyrophosphate, 2mM sodium orthovanadate,

WO 03/100093

PCT/GB03/02307

- 33 -

10mM sodium fluoride, 1mM PMSF, 2 μ g/ml each leupeptin and aprotinin. Lysates were centrifuged for 15minutes at 14,000g, the protein concentration of supernatants was measured using BCA assay reagent (Pierce), and equivalent amounts of protein were separated on 7.5% SDS-PAGE gels and transferred to nitrocellulose. Target protein levels were assessed using antibodies to the β -subunit of the IGF1R or IR (Santa Cruz), phospho-Ser 473 Akt or total Akt (Cell Signalling, New England Biolabs) or β -tubulin (Sigma). Primary antibodies were detected with HRP-conjugated secondary antibodies (Dako), and ECL Plus (Amersham Pharmacia).

15 Results

(1) Down-regulation of IGF1R in MDA-231 cells by antisense oligonucleotides and RNAi:-comparison of the activity of ASOs and RNAi.

20 ASOs and RNAi duplexes were transfected into MDA-231 cells using Cytofectin or Oligofectamine, as described above. After 48hr the cells were lysed, and IGF1R and IR levels were measured by immunoblotting. The intensity of the autoradiographic bands was quantified by densitometry, and IGF1R (or IR) levels were corrected for loading differences. The specific IGF1R or IR results are presented as % IGF1R (or IR) level of that in cells transfected with the same concentration of an appropriate control. This was a scrambled control oligonucleotide for ASOs, and an inverted RNA duplex for RNAi. The results are shown in Table 4 and Figure 2.

35

WO 03/100093

PCT/GB03/02307

- 34 -

Table 4: Effect of IGF1R-ASOs on IGF1R and IR expression in MDA-231 cells

ASO	Protein Level (% control)			
	IGF1R		IR	
	30nM	300nM	30nM	300nM
1	66±23	38±6	112	135
2	84±11	47±5	nd	nd
3	105±18	104±22	nd	nd
4	73±2	49±2	97	92
6	nd	100	nd	100
TSS	141±10	64±19	144	59

- 15 The effects of an ASO and RNAi duplex of equivalent sequence (ASO2 and R2) on IGF1R and IR expression were compared in MDA-231 human breast cancer cells and ME melanoma cells. The transfection protocol was as above, using either Cytofectin (C) or
- 20 Oligofectamine (O; Gibco BRL). After 48hr the cells were lysed and IGF1R and IR levels were determined by immunoblotting. Representative immunoblots are shown in Figure 3.
- a) MDA-231 breast cancer cells transfected with ASO 2 (or scrambled control, Scr) or RNAi (sequence R2 given above) or inverted RNA duplex control (InvRNA) at 20 or 200nM.
- 25 b) MDA-231 cells were transfected with Oligofectamine and RNAi or Inverted control (InvRNA) at 0.1 - 10nM.
- 30 c) ME melanoma cells were transfected with Oligofectamine and RNAi or Inv control duplex at 5 - 500 nM.
- Quantitative analysis of these results is incorporated
- 35 into the graph of ASO effects in MDA-231 (Fig 2).

WO 03/100093

PCT/GB03/02307

- 35 -

It is clear that RNAi causes much more profound inhibition of IGF1R expression than occurs with ASOs. Note that the quantification methods may not be linear at very low protein levels, and it is possible that these analyses underestimate the true extent of IGF1R downregulation. For example the immunoblots (Figure 5) show almost complete inhibition of IGF1R expression (even on these overnight exposures) in MDA-231 breast cancer and ME melanoma cells treated with 10-200nM RNAi.

A second RNAi (R6 in Fig 4; sequence R6 in Table 2) was synthesised corresponding to ASO 6, which hybridizes to IGF1R mRNA on the array with a relative intensity of 0.01. While this does cause downregulation of the IGF1R, the effects were significantly less potent than that of the RNAi corresponding to ASO2 (R2 in Fig 4; sequence R2 in Table 2). RNAi R2 is more effective than RNAi R6 in the following cell lines:

DU145	prostate
A549	NSCLC
UC101	ovary
25 U20S	osteosarcoma
MCF7	breast (ER positive)

This result indicates that intensity of hybridisation to the scanning oligonucleotide array is predictive of the activity of RNAi reagents.

The inhibitory effect of RNAi R2 is partially, though not completely, blocked by the presence of a single base pair mutation (sequence Mut2, in Fig 4), see Figure 4, compare R2 with Mut2. The effect of R6 was less than the effect of the mutant duplex Mut2. This indicates that the efficacy of synthetic 21mer

WO 03/100093

PCT/GB03/02307

- 36 -

RNAi molecules is influenced by secondary structure in, and hence access to, the target region of the mRNA.

- 5 (iv) Comparison of RNAi duplexes R2 and R6 in a range of human and murine cell lines

10 Tumour cells were transfected with 10nM duplexes and IGF1R levels were measured after 48hr. After correction for loading differences IGF1R levels in cells transfected with RNAi were expressed as % of levels in cells transfected with equivalent inverted control duplex.

- 15 Table 4: Effect of R2 and R6 RNAi duplexes in human and murine cell lines

Cell line	Cell type	R2	R6
DU145	Human prostate cancer	34±13	74±8
20 A549	Human non-small cell lung	22±2	102±16
ME	Human melanoma	53±3	104±5
B16	Murine melanoma	45±4	72±14

- 25 In some cell lines eg human ME melanoma and A549 NSCLC, 10nM R6 was essentially inactive. In all cases R2 caused more profound IGF1R downregulation than R6. This suggests that the structural features dictating access are robust and conserved between different cell lines and species.
- 30

(v) Effect of RNAi duplexes of varying length

- 35 Compared with the effects of R2, IGF1R downregulation is less profound after treatment with

WO 03/100093

PCT/GB03/02307

- 37 -

an 18mer duplex (sequence as R2 but lacking the 3 RNA bases at the 3' end). In contrast to the results of Elbashir et al. (2001), a 24mer R2 duplex was observed to be as effective as R2 at 100nM. Results are illustrated in Figure 5. Panel (a) is an immunoblot showing the effect of various RNAis on IGF1R expression at 0.5 and 5nM; panel (b) is a graphical illustration showing the effect of RNAis of varying length, results are presented as % IGF1R level of that in cells transfected with the same concentration of an Inv2 control RNAi. A 27mer R2 duplex was also observed to be as effective as the R2 20mer and 24mer duplexes (data not shown).

15

Example 4-Differential effects of RNAi duplexes on IGF signalling

ME cells were transfected with 100nM RNA duplexes. After 48hr the monolayers were disaggregated and half of each culture was treated with 10nM IGF-1 for 30 min. The cells were lysed and lysates were analysed by immunoblotting for IGF1R, phospho-Ser473-Akt and total Akt (see Figure 6).

25

The IGF1R downregulation that follows transfection with R2 was sufficient to inhibit IGF-I-induced Akt phosphorylation, while R6 caused modest IGF1R downregulation that was not sufficient to block IGF-I signalling to Akt (Figure 6). This is significant because it indicates that R2, which corresponds to an ASO hybridising strongly to the scanning array, not only caused more potent IGF1R downregulation but that this translated to a significant biological advantage; Akt phosphorylation is the major anti-apoptosis pathway downstream of the IGF1R.

35

WO 03/100093

PCT/GB03/02307

- 38 -

Example 5-Growth of cells transfected with RNAi
duplexes in vivo

5 B16 cells were transfected with 200nM RNA
 duplexes (RNA22=R2,22Inv=Inv2) or oligofectamine alone
 (OF) for three consecutive days. On the 4th day cells
 were injected into the flanks of C57BL mice, using
 2.5×10^5 cells/mouse, using groups of 5 mice. Tumours
10 were measured in 2 dimensions every 2-3 days. Tumour
 volumes were calculated as $\pi(a \times b^2)/6$ where a is the
 larger and b the smaller dimension.

 B16 cells transfected with R2 showed reduced
15 growth rate in vivo, (syngeneic C57BL mice) compared
 with cells transfected with inverted control RNA
 duplex (Figure 7). This results indicates that an
 RNAi reagent selected on the basis of hybridisation to
 a scanning array is effective in vivo.

WO 03/100093

PCT/GB03/02307

- 39 -

Claims:

1. A method of preparing an siRNA reagent for use in gene silencing of a target gene by RNA interference, which method comprises:
- 5 (a) preparing a scanning array of antisense oligonucleotides spanning a region of a transcript of the target gene;
- 10 (b) hybridising to the array labelled transcripts of the target gene;
- (c) identifying an oligonucleotide within the array which hybridizes with the labelled transcripts; and
- 15 (d) preparing an siRNA reagent comprising a double-stranded RNA of identical sequence to the oligonucleotide identified in step (c).
2. A method of preparing a pharmaceutical composition comprising an siRNA reagent capable of mediating gene silencing of a target gene by RNA interference, which method comprises:
- 20 preparing an siRNA reagent capable of mediating gene silencing of a target gene by RNA interference according to the method of claim 1 and formulating the siRNA reagent into a pharmaceutical composition
- 25 comprising the siRNA reagent and one or more diluents, excipients or carriers.
3. A method of preparing an expression vector capable of expressing an siRNA reagent for use in gene silencing of a target gene by RNA interference, which method comprises:
- 30 (a) preparing a scanning array of antisense oligonucleotides spanning a region of a transcript of the target gene;
- 35 (b) hybridising to the array labelled transcripts of the target gene;

WO 03/100093

PCT/GB03/02307

- 40 -

(c) identifying an oligonucleotide within the array which hybridizes with the labelled transcripts; and

5 (d) preparing an expression vector capable of expressing an siRNA reagent comprising a double-stranded RNA of identical sequence to the oligonucleotide identified in step (c).

10 4. A method of preparing a pharmaceutical composition comprising an expression vector capable of expressing an siRNA reagent for use in gene silencing of target gene by RNA interference, which method comprises:


15 preparing an expression vector capable of expressing an siRNA reagent for use in gene silencing of a target gene by RNA interference according to the method of claim 3 and formulating the expression vector into a pharmaceutical composition comprising the expression vector and one or more diluents,
20 excipients or carriers.

WO 03/100093

2/9

PCT/GB03/02307

FIG. 1(b).



/images/asohall/pmp-igfr100.gel.clip.rot.rot


Template width: 254.999 Offset: 14.175 First centre: (190, 173)
 Width/offset ratio: 17.9893 Border: 2 Orientation: 0
 Truncated mean: on Aspect ratio: 1000 Array angle: 0
 Scale correction: off Gamma range: (0, 10000)

WO 03/100093

3/9

PCT/GB03/02307




FIG. 2



WO 03/100093

4/9

PCT/GB03/02307

*FIG. 3.***a****b****c**


WO 03/100093

5/9


PCT/GB03/02307

FIG. 4.

(A)



(B)



WO 03/100093

6/9

PCT/GB03/02307






FIG. 5.



WO 03/100093

7/9

PCT/GB03/02307




WO 03/100093

8/9

PCT/GB03/02307

FIG. 7



WO 03/100093

9/9

PCT/GB03/02307

*FIG. 8.***IGF1R cDNA sequence bases 1-1000.**

```
1      TTTT TTTT TTTT TTTGAGAAAGGGAATTT CATCCCAAATAAAAGGAATGAA
51     GTCTGGCTCCGGAGGAGGGTCCCGACCTCGCTGTGGGGGCTCCTGTTTC
101    TCTCCGCCGCGCTCTCGCTCTGGCCGACGAGTGGAGAAATCTGCGGGCCA
151    GGCA TCGACATCCGCAACGACTATCAGCAGCTGAAGCGCCTGGAGAACTG
201    CACGGTGATCGAGGGCTACCTCCACATCCTGCTCATCTCCAAGGCCGAGG
251    ACTACCGCAGCTACCGCTTCCCCAAGCTCACGGTCATTACCGAGTACTTG
301    CTGCTGTTCCGAGTGGCTGGCCTCGAGAGCCTCGGAGACCTCTTCCCCAA
351    CCTCACGGTCATCCGCGGCTGGAAACTCTTCTACAACTACGCCCTGGTCA
401    TCTTCGAGATGACCAATCTCAAGGATATTGGGCTTTACAACCTGAGGACA
451    TTACTCGGGGGGCCATCAGGATTGAGAAAAATGCTGACCTCTGTTACCTC
501    TCCACTGTGGACTGGTCCCTGATCCTGGATGCGGTGTCCTCAATAACTACAT
551    TGTGGGGAATAAGCCCCCAAAGGAATGTGGGGACCTGTGTCCAGGGACCA
601    TGGAGGAGAAGCCGATGTGTGAGAAGACCACCATCAACAATGAGTACAAC
651    TACCGCTGCTGGACCACAAACCGCTGCCAGAAAAATGTGCCCAAGCACGTG
701    TGGGAAGCGGGCGTGCACCGAGAACAATGAGTGCTGCCACCCCGAGTGCC
751    TGGGCAGCTGCAGCGCGCCTGACAAAGACACGGCCTGTGTAGCTTGCCTGC
801    CACTACTACTATGCCGGTGTCTGTGTGCCTGCCTGCCCGCCCAACACCTA
851    CAGGTTTGAGGGCTGGCGCTGTGTGGACCGTGACTTCTGCGCCAACATCC
901    TCAGCGCCGAGAGCAGCGACTCCGAGGGGTTTGTGATCCACGACGGCGAG
951    TGCATGCAGGAGTGCCCCCTCGGGCTTCATCCGCAACGGCAGCCAGAGCAT
```

Underlined are translation start site and region screened by array.

Latter covers 150nt representing bases 536-685 of human IGF1R sequence.

WO 03/100093

PCT/GB03/02307

1

SEQUENCE LISTING

<110> Isis Innovation Limited

<120> Method of selecting targets for gene silencing by RNA interference

<130> SCB/60487/001

<160> 31

<170> PatentIn version 3.1

<210> 1

<211> 18

<212> DNA

<213> artificial sequence

<220>

<223> antisense oligonucleotide

<400> 1

ggcttctctc ccatgggc

18

<210> 2

<211> 20

<212> DNA

<213> Artificial sequence

<220>

<223> Antisense oligonucleotide

<400> 2

aggcttctcc tccatgggcc

20

<210> 3

<211> 20

<212> DNA

<213> Artificial sequence

<220>

<223> Antisense oligonucleotide

<400> 3

gggcttctcc tccaatgtcc

20

<210> 4

<211> 20

<212> DNA

<213> Artificial sequence

<220>

<223> Antisense oligonucleotide

<400> 4

tcttcgcgca cttgctccgc

20

WO 03/100093

PCT/GB03/02307

2

<210> 5
<211> 20
<212> DNA
<213> Artificial sequence

<220>
<223> Antisense oligonucleotide

<400> 5
ctgttcaccg tgcacctgt

20

<210> 6
<211> 18
<212> DNA
<213> Artificial sequence

<220>
<223> Antisense oligonucleotide

<400> 6
ggtcttctca cacatcgg

18

<210> 7
<211> 20
<212> DNA
<213> Artificial sequence

<220>
<223> Antisense oligonucleotide

<400> 7
tggtcttctc acacatcgga

20

<210> 8
<211> 20
<212> DNA
<213> Artificial sequence

<220>
<223> Antisense oligonucleotide

<400> 8
tggtcttctc acacatggga

20

<210> 9
<211> 20
<212> DNA
<213> Artificial sequence

<220>
<223> Antisense oligonucleotide

WO 03/100093

PCT/GB03/02307

3

<400> 9
cctgtacgcg ttgatctcca 20

<210> 10
<211> 20
<212> DNA
<213> Artificial sequence

<220>
<223> Antisense oligonucleotide

<400> 10
agtcgcctag tcgagtcctt 20

<210> 11
<211> 20
<212> DNA
<213> Artificial sequence

<220>
<223> Antisense oligonucleotide

<400> 11
attgttgatg gtggtcttct 20

<210> 12
<211> 20
<212> DNA
<213> Artificial sequence

<220>
<223> Antisense oligonucleotide

<400> 12
tcatgcttgt tgatgtggtt 20

<210> 13
<211> 20
<212> DNA
<213> Artificial sequence

<220>
<223> Antisense oligonucleotide

<400> 13
gcggtagttg tactcattgt 20

<210> 14
<211> 20
<212> DNA
<213> Artificial sequence

<220>

WO 03/100093

PCT/GB03/02307

4

<223> Antisense oligonucleotide

<400> 14

tgtcgttcga gttgatcgta

20

<210> 15

<211> 20

<212> DNA

<213> Artificial sequence

<220>

<223> Antisense oligonucleotide

<400> 15

gtggtccagc agcggtagtt

20

<210> 16

<211> 20

<212> DNA

<213> Artificial sequence

<220>

<223> Antisense oligonucleotide

<400> 16

gacgttagcg tgcgatgtgc

20

<210> 17

<211> 18

<212> DNA

<213> Artificial sequence

<220>

<223> Antisense oligonucleotide

<400> 17

cttattccccc acaatgta

18

<210> 18

<211> 18

<212> DNA

<213> Artificial sequence

<220>

<223> Antisense oligonucleotide

<400> 18

cttgttccccc acaatgta

18

<210> 19

<211> 18

<212> DNA

<213> Artificial sequence

WO 03/100093

PCT/GB03/02307

5

<220>
<223> Antisense oligonucleotide

<400> 19
catagcacat tcttctca

18

<210> 20
<211> 18
<212> DNA
<213> Artificial sequence

<220>
<223> Antisense oligonucleotide

<400> 20
accgttcac agttcact

18

<210> 21
<211> 18
<212> DNA
<213> Artificial sequence

<220>
<223> Antisense oligonucleotide

<400> 21
tcttcggag ccagactt

18

<210> 22
<211> 18
<212> DNA
<213> Artificial sequence

<220>
<223> Antisense oligonucleotide

<400> 22
cagctactcg catgctgc

18

<210> 23
<211> 21
<212> RNA
<213> Artificial sequence

<220>
<223> siRNA reagent

<220>
<221> misc_feature
<222> (20)..(20)
<223> "n" represents "t" thymine

WO 03/100093

PCT/GB03/02307

6

<220>
<221> misc_feature
<222> (21)..(21)
<223> "n" represents "t" thymine

<400> 23
gccgaugugu gagaagaccn n 21

<210> 24
<211> 18
<212> RNA
<213> Artificial sequence

<220>
<223> siRNA reagent

<220>
<221> misc_feature
<222> (17)..(17)
<223> "n" represents "t" thymine

<220>
<221> misc_feature
<222> (18)..(18)
<223> "n" represents "t" thymine

<400> 24
gccgaugugu gagaagann 18

<210> 25
<211> 24
<212> RNA
<213> Artificial sequence

<220>
<223> siRNA reagent

<220>
<221> misc_feature
<222> (23)..(23)
<223> "n" represents "t" thymine

<220>
<221> misc_feature
<222> (24)..(24)
<223> "n" represents "t" thymine

<400> 25
gccgaugugu gagaagacca conn 24

WO 03/100093

PCT/GB03/02307

7

<210> 26
<211> 27
<212> RNA
<213> Artificial sequence

<220>
<223> siRNA reagent

<220>
<221> misc_feature
<222> (26)..(26)
<223> "n" represents "t" thymine

<220>
<221> misc_feature
<222> (27)..(27)
<223> "n" represents "t" thymine

<400> 26
gccgaugugu gagaagacca ccaucnn

27

<210> 27
<211> 21
<212> RNA
<213> Artificial sequence

<220>
<223> siRNA reagent

<220>
<221> misc_feature
<222> (20)..(20)
<223> "n" represents "t" thymine

<220>
<221> misc_feature
<222> (21)..(21)
<223> "n" represents "t" thymine

<400> 27
ccagaagagu guguagccgn n

21

<210> 28
<211> 21
<212> RNA
<213> Artificial sequence

<220>
<223> siRNA reagent

<220>
<221> misc_feature

WO 03/100093

PCT/GB03/02307

8

<222> (20)..(20)
<223> "n" represents "t" thymine

<220>
<221> misc_feature
<222> (21)..(21)
<223> "n" represents "t" thymine

<400> 28
gccgaugugu gugaagaccn n 21

<210> 29
<211> 21
<212> RNA
<213> Artificial sequence

<220>
<223> siRNA reagent

<220>
<221> misc_feature
<222> (20)..(20)
<223> "n" represents "t" thymine

<220>
<221> misc_feature
<222> (21)..(21)
<223> "n" represents "t" thymine

<400> 29
gcccaugugu gagaagaccn n 21

<210> 30
<211> 21
<212> RNA
<213> Artificial sequence

<220>
<223> siRNA reagent

<220>
<221> misc_feature
<222> (20)..(20)
<223> "n" represents "t" thymine

<220>
<221> misc_feature
<222> (21)..(21)
<223> "n" represents "t" thymine

WO 03/100093

PCT/GB03/02307

9

<400> 30
cuacauugug gggauaagn n

21

<210> 31
<211> 21
<212> RNA
<213> Artificial sequence

<220>
<223> siRNA reagent

<220>
<221> misc_feature
<222> (20)..(20)
<223> "n" represents "t" thymine

<220>
<221> misc_feature
<222> (21)..(21)
<223> "n" represents "t" thymine

<400> 31
gaauaagggg uguuacaucn n

21

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☒ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.